

3rd Dose/Booster COVID CDC Guidelines

Pfizer	Moderna
<p>Must wait until 6 months after completion of Pfizer series</p> <ul style="list-style-type: none">• 65+ years old• Residents in Long-Term care settings• 50-64 years old with certain underlying medical conditions• 18-49 years old who are at HIGH risk due to certain underlying medical conditions• 18-64 years old who are at increased risk due to occupational or institutional settings including:<ul style="list-style-type: none">○ First Responders○ Educational Staff○ Food and Agriculture Workers○ Manufacturing Workers○ Corrections Workers○ U.S. Postal Service Workers○ Public Transit Workers○ Grocery Store Workers <p>Underlying medical Conditions:</p> <ul style="list-style-type: none">• Cancer• Chronic kidney disease• Chronic lung disease (COPD, asthma, CF, and pulmonary hypertension)• Dementia or other neurological condition• Diabetes• Down syndrome• Heart conditions (HF, CAD, and hypertension)• HIV infection• Immunocompromised state• Liver disease• Overweight or obesity• Pregnancy• Sickle cell disease• Smoker• Solid blood or organ stem cell transplant• Stroke patient• Substance abuse disorders	<p>3rd dose ONLY applies for immunocompromised patients</p> <ul style="list-style-type: none">• Receiving active cancer treatment• Transplant recipient and currently on medication• Received stem cell transplant within the last 2 years or currently on medication• Moderate/severe immunodeficiency• Advanced/untreated HIV infection• Active high dose corticosteroids (>20 mg prednisone equivalent daily)• Other conditions which cause moderate/severe immunosuppression

Moderna and Johnson & Johnson (Janssen) vaccinations are still being evaluated for booster doses.